



# Efficient Liquid Media for Encystation of Pathogenic Free-Living Amoebae

Hae-Jin Sohn<sup>1</sup>, Heekyoung Kang<sup>1</sup>, Ga-Eun Seo<sup>1</sup>, Jong-Hyun Kim<sup>2</sup>, Suk-Yul Jung<sup>3</sup>, Ho-Joon Shin<sup>1,\*</sup>

<sup>1</sup>Department of Microbiology, Ajou University School of Medicine, and Department of Biomedical Science, Graduate School of Ajou University, Suwon 16499, Korea; <sup>2</sup>Institute of Animal Medicine, College of Veterinary Medicine, Gyeongsang National University, Jinju 52828, Korea; <sup>3</sup>Department of Biomedical Laboratory Science, Molecular Diagnostics Research Institute, School of Health and Medicine, Namseoul University, Cheonan 31020, Korea

**Abstract:** Pathogenic *Naegleria fowleri*, *Acanthamoeba castellanii*, and *Acanthamoeba polyphaga*, are distributed worldwide. They are causative agents of primary amoebic meningoencephalitis or acanthamoebic keratitis in humans, respectively. Trophozoites encyst in unfavorable environments, such as exhausted food supply and desiccation. Until recently, the method of *N. fowleri* encystation used solid non-nutrient agar medium supplemented with heat-inactivated *Escherichia coli*; however, for the amoebic encystment of *Acanthamoeba* spp., a defined, slightly modified liquid media is used. In this study, in order to generate pure *N. fowleri* cysts, a liquid encystment medium (buffer 1) modified from Page's amoeba saline was applied for encystation of *N. fowleri*. *N. fowleri* cysts were well induced after 24 hr with the above defined liquid encystment medium (buffer 1). This was confirmed by observation of a high expression of differential mRNA of *nfa1* and actin genes in trophozoites. Thus, this liquid medium can replace the earlier non-nutrient agar medium for obtaining pure *N. fowleri* cysts. In addition, for cyst formation of *Acanthamoeba* spp., buffer 2 (adjusted to pH 9.0) was the more efficient medium. To summarize, these liquid encystment media may be useful for further studies which require axenic and pure amoebic cysts.

**Key words:** *Naegleria fowleri*, *Acanthamoeba castellanii*, *Acanthamoeba polyphaga*, encystation, encystment medium

## INTRODUCTION

Pathogenic free-living amoebae, *Naegleria fowleri*, *Acanthamoeba castellanii*, and *Acanthamoeba polyphaga*, are distributed worldwide, and are the known causative agents of primary amoebic meningoencephalitis (PAM) or acanthamoebic keratitis (AK) in humans, respectively [1]. PAM cases are associated with recreational activities in freshwater, whereas AK cases have become a concern in people wearing contact lenses. The ubiquitous distribution in nature of these free-living amoebae shows 2 distinct forms in their life-cycle: trophozoites and cysts (sometimes flagellate in *Naegleria*). In unfavorable environments, such as exhausted food supply, desiccation, and low temperature, amoebic trophozoites are known to undergo encystation. The cysts are characterized by a round shape and the

presence of 2 cyst walls (endo- and ectocyst), especially distinct in the wrinkled ectocyst in *Acanthamoeba* spp. [2]. The encystation is an important event for the survival, resistance, and ability of disease induction of these free-living amoebae [3-6].

Until recently, a useful method of *N. fowleri* encystation was the use of solid non-nutrient agar medium supplied with heat-inactivated *Escherichia coli*. The solid medium has a limitation for cultivation and harvesting axenic and massive amoebic cysts. Current studies in *Naegleria* infection indicate that the encystation of *N. fowleri* trophozoites is required to study the pathophysiology, pathogenic mechanisms, immunology, and genomics. On the other hand, since a defined liquid media for the encystment of *Acanthamoeba* spp. has been established [6], many researchers have lately applied acanthamoebic encystation with a little modification [7-10].

To obtain a large amount of pure *N. fowleri* cysts (or precysts), a liquid encystment medium modified from Page's amoeba saline [1] was applied on encystation of *N. fowleri* trophozoites. In addition, the efficient usage of 2 liquid encystment media generally used for the cyst formation of *Acanthamoeba* spp. [6,11] was compared.

•Received 15 February 2017, revised 4 April 2017, accepted 10 April 2017.

\*Corresponding author (hjshin@ajou.ac.kr)

© 2017, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## MATERIALS AND METHODS

### Free-living amoebae cultivation

Trophozoites of *N. fowleri* (Cater NF69 strain, ATCC No. 30215, Manassas, Virginia, USA) were axenically cultured at 37°C in Nelson's medium containing 10% fetal bovine serum [12], and *A. castellanii* and *A. polyphaga* were axenically cultured at 30°C in PYG medium [13].

### Encystation of free-living amoebae

Cysts or precysts (did not fully showed the 2-distinct cyst walls) of *N. fowleri*, *A. castellanii*, and *A. polyphaga* were induced by cultivating on 3 kinds of encystment media (buffer 1: 120 mM NaCl, 0.03 mM MgCl<sub>2</sub>, 1 mM NaHPO<sub>4</sub>, 1 mM KH<sub>2</sub>PO<sub>4</sub>, 0.03 mM CaCl<sub>2</sub>, 0.02 mM FeCl<sub>2</sub>, H 6.8; buffer 2 [11]: 95 mM NaCl, 5 mM KCl, 8 mM MgSO<sub>4</sub>, 0.4 mM CaCl<sub>2</sub>, 1 mM NaHCO<sub>3</sub>, 20 mM Tris-HCl, pH 9.0; buffer 3 [6]: 100 mM KCl, 8 mM MgSO<sub>4</sub>, 0.4 mM CaCl<sub>2</sub>, 20 mM 2-amino-2-methyl-1,3-propanediol, pH 7.6) (Table 1). Amoebic trophozoites (approximately 2 × 10<sup>6</sup> cells) were washed with PBS (pH 7.4) twice, and incubated in 24-well plates with 5 ml of each medi-

**Table 1.** Compositions of encystation media for *Naegleria* (buffer 1), *Acanthamoeba* spp. (buffers 2 and 3), and Page's amoeba saline

Buffer 1. <i>Naegleria</i> Encystment media (pH6.8)	
120 mM	NaCl
0.03 mM	MgCl <sub>4</sub>
1 mM	Na <sub>2</sub> HPO <sub>4</sub>
1 mM	KH <sub>2</sub> PO <sub>4</sub>
0.03 mM	CaCl <sub>2</sub>
0.02 mM	FeCl <sub>2</sub>
Buffer 2. <i>Acanthamoeba</i> Encystment media (pH9.0)	
95 mM	NaCl
5 mM	KCl
8 mM	MgSO <sub>4</sub>
1 mM	NaHCO <sub>3</sub>
0.4 mM	CaCl <sub>2</sub>
20 mM	Tris-Cl (Ph9.0)
Buffer 3. <i>Acanthamoeba</i> Encystment media (pH7.6)	
0.1 mM	KCl
8 mM	MgSO <sub>4</sub>
0.4 mM	CaCl <sub>2</sub>
20 mM	2-amino-2-methyl-1, 3-propanediol
Page's amoeba saline (pH6.8-7.9)	
2 mM	NaCl
0.03 mM	MgSO <sub>4</sub>
0.03 mM	CaCl <sub>2</sub>
1 mM	Na <sub>2</sub> HPO <sub>4</sub>
1 mM	KH <sub>2</sub> PO <sub>4</sub>

um, at 30°C or 37°C. Using an optical microscope (Olympus, Shinjuku, Tokyo, Japan), the morphological changes were observed after encystation, and final cysts were re-cultured with fresh Nelson or PYG media, in order to observe the recovered trophozoites.

### Differential mRNA expression from trophozoites and cysts of free-living amoebae


Total RNA from trophozoites and cysts (or precysts) of *N. fowleri*, *A. castellanii*, and *A. polyphaga* were prepared using an isolation kit Rnasymini kit (QIAGEN, Hilden, Germany); cDNAs were synthesized from 10 µg of total RNA using cDNA Synthesis kit (Invitrogen, Carlsbad, California, USA). RT-PCR was performed using specific primer for *nfa1* and *nf-actin* genes for *Naegleria*, and *actin* and *atg8* genes for *Acanthamoeba* [5,14, 15]. The reference genes were the *p3* gene (*N. fowleri*-specific chromosomal DNA sequence; forward 5'GCTATCGAATG-GATTCAAGC and reverse 5'CACTACTCGTGGAAGGCTTA) and *18S rRNA* gene [5]. PCR condition was 95°C for 5 min, 30 cycles at 95°C for 1 min, 50-55°C for 1 min, 72°C for 1 min, and final extension for 10 min at 72°C.

## RESULTS


### Morphological observation of encysted amoebae

Trophozoites of *N. fowleri* cultured with buffer 1 showed the rounding morphological change at 6 hr, and precysts and cysts were observed at 24 hr. After 48 hr cultivation, a large proportion of *N. fowleri* trophozoites were encysted (Fig. 1). On culturing the *N. fowleri* trophozoites in buffer 2, the morphological changes (such as rounding), precysts and cysts were induced later than seen in buffer 1 (Fig. 1). On all occasions, the induced precysts and cysts at 48 hr were recovered into active trophozoites in fresh Nelson's medium (Fig. 1). As a control group, *N. fowleri* trophozoites cultured in distilled water (DW) did not encyst fully. In addition, when they were incubated with staurosporine, a reagent known to induce cell-death, *N. fowleri* trophozoites transformed to their round forms, but live trophozoites could not be recovered in Nelson's medium (Fig. 1).


Trophozoites *A. castellanii* cultured with buffer 2 and 3 showed rounding at day 1, and precysts and cysts were well observed at day 2. After cultivation for 3 days, a large proportion of *A. castellanii* trophozoites were encysted (Fig. 2). Culturing *A. castellanii* trophozoites in buffer 2, the induction of precysts and cysts was better than those of buffer 3 (Fig. 2). On




**Fig. 1.** Morphological changes of *N. fowleri* trophozoites into precysts (or cysts) cultured with encystation media for 6, 24, and 48 hr. Buffer 1 is an effective medium. Staurosporine induced rounding, but trophozoites were not recovered in Nelson's medium.



**Fig. 2.** Morphological changes of *A. castellanii* trophozoites into precysts (or cysts) cultured in encystation media for 1, 2, and 3 days. The effective medium is buffer 2. Cysts recovered to trophozoites in PYG medium. Arrows indicate mature cysts.



**Fig. 3.** Morphological changes of *A. polyphaga* trophozoites into precysts (or cysts) cultured in encystation media for 1, 2, and 3 days. Buffer 2 is an effective medium. Cysts recovered to trophozoites in PYG medium.




**Fig. 4.** The mRNA expressions of *nfa1* and *nf-actin* genes on *N. fowleri* trophozoites (Troph), precysts, and cysts (PreC) at 48 hr post cultivation with buffer 1. The *p3* mRNA expression is used as the control.

all occasions, induced precysts and cysts for 3 days recovered into active trophozoites in fresh PYG medium (Fig. 2). Furthermore, *A. castellanii* trophozoites cultured with buffer 1 (shown the well encystation in the case of *N. fowleri*) did not encyst fully (Fig. 2). When encystation was continued for 7 days, similar results were observed, although the number of cysts decreased in accordance with incubation periods (data not shown). In addition, in the case of *A. polyphaga*, the encystation results were similar to those of *A. castellanii* (Fig. 3).

#### Differential mRNA expression in trophozoites and cysts of amoebae

In the results of differential mRNA expression between trophozoites and cysts (precyst) of *N. fowleri*, the *nfa1* gene was highly expressed in the trophozoites (Fig. 4). In addition, the *nf-actin* gene on trophozoites was also more expressed than in cysts (Fig. 4). In the differential mRNA expression of acanthamoebic actin gene, over-expression was observed in trophozoites of *A. castellanii* and *A. polyphaga* (Fig. 5). In contrast,



**Fig. 5.** The mRNA expressions of *actin* and *atg8* genes on *A. castellanii* (A) and *A. polyphaga* (B) trophozoites, precysts and cysts. The 18S rRNA mRNA expression is used as the control.

the *atg8* gene which is associated with cyst formation in *Acanthamoeba* spp. [11] was expressed more in cysts (Fig. 5).

## DISCUSSION

*N. fowleri*, known as “brain-eating amoeba”, causes acute PAM in children and young adults with a history of diving in freshwater and recreational activities in amoebae-contaminated water in the hot summer [2,16]. In these latter days, death due to PAM is rearing its ugly head as a Neglected Tropical Disease (NTD) in countries of Asia, such as India, Pakistan, Taiwan, Thailand, and Vietnam [17]. In addition, the AK human case from contact lens wearers has been increasing worldwide [18].

In recent many studies on free-living amoebae, such as *N. fowleri* and *A. castellanii*, the encystation of amoebic trophozoite is required to study the pathophysiology, pathogenic mechanisms, immunology, and genomics. For the amoebic encystation of *Acanthamoeba* spp., in previous studies, a defined and slightly modified liquid media has been used [6,9,10]. In *N. fowleri* encystation, however, the solid non-nutrient agar medium supplemented with heat-inactivated *E. coli* has been used, in spite of the possible contamination of *E. coli* proteins and smaller numbers of harvested cysts [4,19,20].

In this study, to obtain a large amount of pure *N. fowleri* cysts (or precysts), a liquid encystment medium modified from

Page’s amoeba saline (named as the encystment buffer 1) was applied on *N. fowleri* encystation. As the results of *N. fowleri* encystation with buffer 1, amoebic cysts were well induced at 48 hr post incubation. And then, the induced cysts at 48 hr were well recovered into active trophozoites in fresh Nelson’s medium. In addition, the above morphological encystation was confirmed by observing the differential mRNA expression of the pseudopodia-specific *nfa1* gene [14] and the *nf-actin* gene [15] between trophozoites and cysts of *N. fowleri*. On the present results, the *nfa1* and *nf-actin* gene on trophozoites was also more expressed than in cysts. These results proved the encystation of *N. fowleri* trophozoites. Thus, the present method used a liquid encystment medium may replace the previous method of *N. fowleri* encystation used a solid non-nutrient agar medium with heat-inactivated *E. coli*.

On the morphological findings of *Acanthamoeba* encystation, buffer 2 (modified to pH 9.0) was also a more effective agent in this study, which was mainly used for many studies [6,8,9,11]. In addition, the present results were confirmed by the differential mRNA expression between trophozoites and cysts. The over-expression of *actin* gene was observed in trophozoites of *A. castellanii* and *A. polyphaga*. In contrast, the *atg8* gene was expressed more in cysts, as which is associated with cyst formation in *Acanthamoeba* spp. [11]. The above results indicate that buffer 2 is better for the encystation of *A. castellanii* and *A. polyphaga* trophozoites.

Finally, 2 liquid encystment media, buffer 1 for *Naegleria* and buffer 2 for *Acanthamoeba* may be useful for further studies which require axenic and pure amoebic cysts.

## ACKNOWLEDGMENT

This research was supported by a Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, the Republic of Korea (no. 2015R1D1A1A01056862).

## CONFLICT OF INTEREST

We have no conflict of interest related to this study.

## REFERENCES

- Ma P, Visvesvara GS, Martinez AJ, Theodore FH, Daggett PM, Sawyer TK. *Naegleria* and *Acanthamoeba* infections: review. *Rev Infect Dis* 1990; 12: 490-513.
- Schuster FL, Visvesvara GS. Free-living amoebae as opportunistic and non-opportunistic pathogens of humans and animals. *Int J Parasitol* 2004; 34: 1001-1027.
- Zhang L, Marciano-Cabral F, Bradley SG. Effects of cyclophosphamide and a metabolite, acrolein, on *Naegleria fowleri* in vitro and in vivo. *Antimicrob Agents Chemother* 1988; 32: 962-965.
- Visvesvara GS, Moura H, Schuster FL. Pathogenic and opportunistic free-living amoebae: *Acanthamoeba* spp., *Balamuthia mandrillaris*, *Naegleria fowleri*, and *Sappinia diploidea*. *FEMS Immunol Med Microbiol* 2007; 50: 1-26.
- Moon EK, Chung DI, Hong YC, Kong HH. Autophagy protein 8 mediating autophagosome in encysting *Acanthamoeba*. *Mol Biochem Parasitol* 2009; 168: 43-48.
- Neff RJ, Benton WF, Wilborn M. Induction of synchronous encystment (differentiation) in *Acanthamoeba* sp. *Methods Cell Biol* 1964; 1:55-83.
- Bowers B, Korn ED. The fine structure of *Acanthamoeba castellanii* (Neff strain). II. Encystment. *J Cell Biol* 1969; 41: 786-805.
- Chagla AH, Griffiths AJ. Growth and encystation of *Acanthamoeba castellanii*. *J Gen Microbiol* 1974; 85: 139-145.
- Moon EK, Park HR, Quan FS, Kong HH. Efficacy of Korean multipurpose contact lens disinfecting solutions against *Acanthamoeba castellanii*. *Korean J Parasitol* 2016; 54: 697-702.
- Abedkhozasteh H, Niyati M, Rezaei S, Mohebbi M, Farnia S, Kazemi-Rad E, Roozafzoon R, Sianati H, Rezaeian M, Heidari M. Identifying differentially expressed genes in trophozoites and cysts of *Acanthamoeba* T4 genotype: Implications for developing new treatments for *Acanthamoeba* keratitis. *Eur J Protistol* 2015; 51: 34-41.
- Moon EK, Chung DI, Hong YC, Ahn TI, Kong HH. *Acanthamoeba castellanii*: gene profile of encystation by ESTs analysis and KOG assignment. *Exp Parasitol* 2008; 119: 111-116.
- Willaert E. Isolation and in vitro culture of the amoeba of the genus *Naegleria*. *Ann Soc Belges Med Trop Parasitol Mycol* 1971; 51: 701-708.
- Visvesvara GS, Balamuth W. Comparative studies on related free-living and pathogenic amoebae with special reference to *Acanthamoeba*. *J Protozool* 1975; 22: 245-256.
- Shin HJ, Cho MS, Jung SU, Kim HI, Park S, Kim HJ, Im KI. Molecular cloning and characterization of a gene encoding a 13.1 kDa antigenic protein of *Naegleria fowleri*. *J Eukaryot Microbiol* 2001; 48: 713-717.
- Sohn HJ, Kim JH, Shin MH, Song KJ, Shin HJ. The *Nf-actin* gene is an important factor for food-cup formation and cytotoxicity of pathogenic *Naegleria fowleri*. *Parasitol Res* 2010; 106: 917-924.
- Kang H, Seong GS, Sohn HJ, Kim JH, Lee SE, Park MY, Lee WJ, Shin HJ. Effective PCR-based detection of *Naegleria fowleri* from cultured sample and PAM-developed mouse. *Eur J Protistol* 2015; 51:401-408.
- Siddiqui R, Khan NA. Primary amoebic meningoencephalitis caused by *Naegleria fowleri*: an old enemy presenting new challenges. *PLoS Negl Trop Dis* 2014;8: e3017.
- Lorenzo-Morales J, Khan NA, Walochnik J. An update on *Acanthamoeba* keratitis: diagnosis, pathogenesis and treatment. *Parasite* 2015; 22: 10.
- Visvesvara GS, De Jonckheere JE, Marciano-Cabral F, Schuster FL. Morphologic and molecular identification of *Naegleria dunnebackei* n. sp. isolated from a water sample. *J Eukaryot Microbiol* 2005; 52: 523-531.
- Chávez-Munguía B, Segovia-Gamboa N, Salazar-Villatoro L, Omaña-Molina M, Espinosa-Cantellano M, Martínez-Palomo A. *Naegleria fowleri*: enolase is expressed during cyst differentiation. *J Eukaryot Microbiol* 2011; 58: 463-468.